

ON THE c -DOMINATING ESTRADA INDEX OF A GRAPH

AKBAR JAHANBANI AND HAJAR SHOOSHTARI

ABSTRACT. Let G be a graph with n vertices then the c -dominating matrix of G is the square matrix of order n whose (i, j) -entry is equal to 1 if the i -th and j -th vertex of G are adjacent, is also equal to 1 if the $i = j, v_i \in D_c$ and zero otherwise. The c -dominating energy of a graph G , is defined as the sum of the absolute values of all eigenvalues the c -dominating matrix. The main purposes of this paper are to introduce the c -dominating Estrada index of a graph. Moreover, obtain upper and lower bounds for the c -dominating Estrada index and investigate the relations between the c -dominating Estrada index and the c -dominating energy.

1. INTRODUCTION

Let G be such a graph with vertex set $V(G)$ and edge set $E(G)$. The degree of a vertex $v \in V(G)$ is the number of vertices adjacent to v and is denoted by $d_G(v)$. Let the vertices of G be labeled as v_1, v_2, \dots, v_n . The vertex u and v are adjacent if $uv \in E(G)$. The open(closed) neighborhood of a vertex $v \in V(G)$ is $N(v) = \{u : uv \in E(G)\}$ and $N[v] = N(v) \cup v$ respectively. The degree of a vertex $v \in V(G)$ is denoted by $d_G(v)$ and is defined as $d_G(v) = |N(v)|$. A vertex $v \in V(G)$ is pendant if $|N(v)| = 1$ and is called support vertex if it is adjacent to pendant vertex. Let us denote the complete graph, and the path having n vertices by K_n and P_n , respectively. The complement of G , denoted by \overline{G} , is a graph which has the same vertices as G , and in which two vertices are adjacent if and only if they are not adjacent in G .

Any vertex $v \in V(G)$ with $|N(v)| > 1$ is called internal vertex. A subset $D \subseteq V(G)$ is called dominating set if $N[D] = V(G)$. The minimum cardinality of such a set D is called the domination number $\gamma(G)$ of G . A dominating set D is connected if the subgraph induced by D is connected. The minimum cardinality of connected dominating set D is called the connected dominating number $\gamma_c(G)$ of G [28]. The *adjacency* matrix of a graph G is a square matrix $A(G) = [a_{ij}]$ of order n , defined via

$$a_{ij} = \begin{cases} 1 & \text{if the vertices } v_i \text{ and } v_j \text{ of } G \text{ are adjacent} \\ 0 & \text{otherwise.} \end{cases}$$

The *eigenvalues* of $A(G)$ are the adjacency eigenvalues of G , they are labeled as $\lambda_1, \lambda_2, \dots, \lambda_n$. The graph energy is a graph-spectrum-based quantity, introduced in the 1970s.

2020 *Mathematics Subject Classification.* 05C50.

Key words and phrases. Eigenvalue of graph, Energy, c -dominating Estrada index, c -dominating Energy, Connected dominating number.

After a latent period of 20–30 years, it became a popular topic of research both in mathematical chemistry and in “pure” spectral graph theory, resulting in over 600 published papers. The first paper in which graph energy was defined as the sum of absolute values of the eigenvalues of the $(0; 1)$ -adjacency matrix, namely as

$$(1) \quad E = E(G) = \sum_{i=1}^n |\lambda_i|.$$

The concept of energy first by *I. Gutman* be used to approximate the total π -electron energy of a *molecule* (see, e.g. [9, 10]). For more details on energy mathematical properties see, ([1, 3], [16]-[19]).

A graph-spectrum-based graph invariant, recently put forward by Estrada [7, 8], is defined as

$$EE = EE(G) = \sum_{i=1}^n e^{\lambda_i}.$$

EE is nowadays usually referred to as the Estrada index, see [26]. Although invented only a few years ago [7], the Estrada index has already found numerous applications. It was used to quantify the degree of folding of long-chain molecules, especially proteins [8]. Some mathematical properties of the Estrada index were established in [4, 13, 14, 15]. One of most important properties is the following:

$$(2) \quad EE = \sum_{k=1}^{\infty} \frac{M_k(G)}{k!}.$$

Denoting by $M_k = M_k(G)$ to the k -th moment of the graph G , we get where, $M_k = M_k(G) = \sum_{i=1}^n (\lambda_i)^k$. It is well known that $M_k(G)$ is equal to the number of closed walks of length k in G .

Refer to references [20, 21, 22, 23, 24, 27] to see recent work on the Estrada index.

The c -dominating matrix of G denoted by $D_c(G) = (d_{ij})$, defined in [11] as following

$$d_{ij} = \begin{cases} 1 & \text{if } i = j \text{ and } v_i \in D_c, \\ 1 & \text{if the vertices } v_i \text{ and } v_j \text{ of } G \text{ are adjacent,} \\ 0 & \text{otherwise.} \end{cases}$$

The eigenvalues of $D_c(G)$ labeled as $\theta_1 \geq \theta_2 \geq \dots \geq \theta_n$ are said to be D_c -eigenvalues of G and their collection is called the c -dominating spectrum or D_c -spectrum of G . The spectral radius of G , denoted by $\theta_1(G)$, is the largest eigenvalue of $D_c(G)$.

In [11], the c -dominating energy of G is then defined as following

$$E_c = E_c(G) = \sum_{i=1}^n |\theta_i|.$$

Similarly to the definition of Estrada index, we define the c -dominating Estrada index of G , as following

$$EE_c = EE_c(G) = \sum_{i=1}^n e^{\theta_i}.$$

Now, here, we denoted

$$N_k = \sum_{i=1}^n (\theta_i)^k$$

hence, with an argument similar to (2), we can write that

$$EE_c(G) = \sum_{i=1}^{\infty} \frac{N_k}{k!}.$$

This paper is organized as follows. In Section 2, we give a list of some previously known results. In Section 3, introducing c -dominating Estrada index and we establish upper and lower bounds for it. In Section 4, we investigate the relations between c -dominating Estrada index and c -dominating energy. Note that all graphs considered in this paper are simple.

2. PRELIMINARIES AND KNOWN RESULTS

In this section, we shall list some previously known results that will be needed in the next sections. Now we give some lemmas which will be needed later. The following result comes from [11].

Lemma 1. [11] *Let G be a graph with n vertices and c -dominating matrix D_c . Then*

$$(3) \quad (1) \quad N_1 = \sum_{i=1}^n \theta_i = \text{tr}(D_c) = \gamma_c(G) = \gamma_c,$$

$$(4) \quad (2) \quad N_2 = \sum_{i=1}^n (\theta_i)^2 = \text{tr}(D_c^2) = 2m + \gamma_c(G).$$

Now, we obtain upper bound for N_4 that will be need to obtain the result.

Lemma 2. *Let G be a graph with n vertices and $\theta_1 \geq \theta_2 \geq \dots \geq \theta_n$ be a non-increasing arrangement of eigenvalues of G . Then, the following inequality is valid*

$$(5) \quad \sum_{i=1}^n \theta_i^4 \leq (\theta_n^2 + \theta_1^2) (2m + \gamma_c(G)) - n\theta_1^2\theta_n^2.$$

Proof. Let a_1, a_2, \dots, a_n and b_1, b_2, \dots, b_n be real numbers for which there exist real constants R and r , so that for each $i, i = 1, 2, \dots, n$, holds $ra_i \leq b_i \leq Ra_i$. Then the following inequality is valid (see [5])

$$(6) \quad \sum_{i=1}^n b_i^2 + rR \sum_{i=1}^n a_i^2 \leq (r + R) \sum_{i=1}^n a_i b_i.$$

Equality in (6) holds if and only if for at least one $i, 1 \leq i \leq n$ holds $ra_i = bi = Ra_i$. Setting $a_i := 1, b_i := \theta_i^2, r := \theta_n^2$ and $R := \theta_1^2, i = 1, 2, \dots, n$, in Inequality (6), we get

$$\sum_{i=1}^n \theta_i^4 + \theta_1^2 \theta_n^2 \sum_{i=1}^n 1 \leq (\theta_n^2 + \theta_1^2) \sum_{i=1}^n \theta_i^2.$$

By Equality (4), we have,

$$\sum_{i=1}^n \theta_i^2 = 2m + \gamma_c(G).$$

Also, we know that

$$\sum_{i=1}^n 1 = n,$$

Therefore, the above inequality becomes

$$\sum_{i=1}^n \theta_i^4 \leq (\theta_n^2 + \theta_1^2) (2m + \gamma_c(G)) - n\theta_1^2 \theta_n^2.$$

□

Remark 1. For any real x , the power-series expansion of e^x , is the following

$$(7) \quad e^x = \sum_{k \geq 0} \frac{x^k}{k!}.$$

Remark 2. For nonnegative x_1, x_2, \dots, x_n and $k \geq 2$,

$$(8) \quad \sum_{i=1}^n (x_i)^k \leq \left(\sum_{i=1}^n x_i^2 \right)^{\frac{k}{2}}.$$

Lemma 3. (Rayleigh-Ritz) [12] If B is a real symmetric $n \times n$ matrix with eigenvalues $\lambda_1(B) \geq \lambda_2(B) \geq \dots \geq \lambda_n(B)$, then for any $X \in \mathbb{R}^n, (X \neq 0)$,

$$X^t B X \leq \lambda_1(B) X^t X.$$

Equality holds if and only if X is an eigenvector of B , corresponding to the largest eigenvalue $\lambda_1(B)$.

3. BOUNDS FOR THE c -DOMINATING ESTRADA INDEX OF GRAPHS

In this section, we consider the c -dominating Estrada index of graph G . Also we present lower and upper bounds for the c -dominating Estrada index in terms of the number of vertices, edges and the connected dominating number.

At first, we state the following useful lemma.

Lemma 4. Let G be a connected graph with $n \geq 2$ vertices. Then the spectral radius of the c -dominating matrix is bounded from below as

$$(9) \quad \theta_1 \geq \frac{2D_c(G)}{n}.$$

Proof. Let $D_c = ||d_{ij}||$ be c -dominating matrix corresponding to D_c . By Lemma 3, for any vector $X = (x_1, x_2, \dots, x_n)^t$,

$$\begin{aligned} X^t D_c X &= \left(\sum_{j,j \sim 1}^n x_j d_{j1}, \sum_{j,j \sim 2}^n x_j d_{j2}, \dots, \sum_{j,j \sim n}^n x_j d_{jn} \right)^t X \\ (10) \qquad &= 2 \sum_{i \sim j} d_{ij} x_i x_j \end{aligned}$$

because $d_{ij} = d_{ji}$. Also,

$$(11) \qquad X^t X = \sum_{i=1}^n x_i^2.$$

Using Eqs. (10) and (11), by Lemma 3, we obtain

$$(12) \qquad \theta_1 \geq \frac{2 \sum_{i \sim j} d_{ij} x_i x_j}{\sum_{i=1}^n x_i^2}.$$

Since (12) is true for any vector X , by putting $X = (1, 1, \dots, 1)^t$, we have

$$\theta_1 \geq \frac{2D_c(G)}{n}.$$

□

We are now ready to give some new bounds for $EE_c(G)$.

Theorem 1. *Let G be a graph with n vertices. Then*

$$EE_c(G) \geq e^{\frac{2D_c(G)}{n}} + \frac{n-1}{e^{\left(\frac{\frac{2D_c(G)-\gamma_c}{n}}{n-1}\right)}}.$$

Proof. By the definition of c -dominating Estrada index and using arithmetic-geometric mean inequality, we obtain

$$\begin{aligned} EE_c(G) &= e^{\theta_1} + e^{\theta_2} + \dots + e^{\theta_n} \\ (13) \qquad &\geq e^{\theta_1} + (n-1) \left(\prod_{i=2}^n e^{\theta_i} \right)^{\frac{1}{n-1}} \end{aligned}$$

$$(14) \qquad = e^{\theta_1} + (n-1) \left(e^{\gamma_c - \theta_1} \right)^{\frac{1}{n-1}} \quad \text{by Equality (3).}$$

Now we consider the following function

$$f(x) = e^x + \frac{n-1}{e^{\frac{x-\gamma_c}{n-1}}}$$

for $x > 0$. We have

$$f(x) \geq e^x + \frac{n-1}{e^{\frac{x-\gamma_c}{n-1}}}$$

It is easy to see that f is an increasing function for $x > 0$. From the Equation (14) and Lemma 4, we get

$$EE_c(G) \geq e^{\frac{2D_c(G)}{n}} + \frac{n-1}{e^{\left(\frac{2D_c(G)-\gamma_c}{n-1}\right)}}.$$

□

Theorem 2. Let G be a graph with $n \geq 5$ vertices, $m \geq 5$ edges and $\gamma_c(G)$ the connected dominating number. Then

$$(15) \quad EE_c(G) \leq n-1 + e^{\sqrt{2m+\gamma_c(G)-1}}.$$

Proof. Let the number of positive eigenvalues of G be n_+ . Since $f(x) = e^x$ monotonically increases in the interval $(-\infty, +\infty)$ and $m \neq 0$, we get

$$(16) \quad \begin{aligned} EE_c(G) &= \sum_{i=1}^n e^{\theta_i} < n - n_+ + \sum_{i=1}^{n_+} e^{\theta_i} = n - n_+ + \sum_{i=1}^{n_+} \sum_{k \geq 0} \frac{(\theta_i)^k}{k!} \\ &= n + \sum_{k \geq 1} \frac{1}{k!} \sum_{i=1}^{n_+} (\theta_i)^k \\ &\leq n + \sum_{k \geq 1} \frac{1}{k!} \left[\sum_{i=1}^{n_+} \theta_i^2 \right]^{\frac{k}{2}} \\ &= n + \sum_{k \geq 1} \frac{1}{k!} \left[2m + \gamma_c(G) - \sum_{i=n_++1}^n \theta_i^2 \right]^{\frac{k}{2}}. \end{aligned}$$

Since every graph with $n \geq 5$ vertices and $m \geq 5$ edges has P_5 as its induced sub-graph and by direct calculation it can be easily seen that the spectrum of P_5 are $\theta_1 = 2.618, \theta_2 = 1.618, \theta_3 = 0.382, \theta_4 = -1.000, \theta_5 = -0.618$, it follows from the interlacing inequalities that $\theta_n \leq 1$, which implies that, $\sum_{i=n_++1}^n \theta_i^2 \geq 1$. Consequently,

$$EE_c(G) \leq n + \sum_{k \geq 1} \frac{1}{k!} \left[2m + \gamma_c(G) - 1 \right]^{\frac{k}{2}} = n - 1 + e^{\sqrt{2m+\gamma_c(G)-1}}.$$

□

Theorem 3. Let G be a graph with n vertices, m edges and $\gamma_c(G)$ the connected dominating number. Then

$$EE_c(G) \leq n - 1 + \gamma_c + \frac{2m + \gamma_c}{2} + \frac{N_3}{6} - N_4^{1/4} - \frac{N_4^{1/2}}{2} - \frac{N_4^{3/4}}{6} + e^{\sqrt[4]{2m+\gamma_c(G)-n\theta_1\theta_n}}.$$

Proof. By definition of the c -dominating Estrada index, we have

$$\begin{aligned}
 EE_c(G) &= \sum_{i=1}^n e^{\theta_i} = \sum_{i=1}^n \sum_{k=0}^{\infty} \frac{\theta_i^k}{k!} = n + \sum_{i=1}^n \theta_i + \frac{1}{2} \sum_{i=1}^n \theta_i^2 + \frac{1}{6} \sum_{i=1}^n \theta_i^3 + \sum_{i=1}^n \sum_{k=4}^{\infty} \frac{\theta_i^k}{k!} \\
 &= n + \sum_{i=1}^n \theta_i + \frac{1}{2} \sum_{i=1}^n \theta_i^2 + \frac{1}{6} \sum_{i=1}^n \theta_i^3 + \sum_{i=1}^n \sum_{k=4}^{\infty} \frac{|\theta_i|^k}{k!} \\
 &= n + \sum_{i=1}^n \theta_i + \frac{1}{2} \sum_{i=1}^n \theta_i^2 + \frac{1}{6} \sum_{i=1}^n \theta_i^3 + \sum_{k=4}^{\infty} \frac{1}{k!} \sum_{i=1}^n (\theta_i^k)^{\frac{k}{4}} \\
 &\leq n + \sum_{i=1}^n \theta_i + \frac{1}{2} \sum_{i=1}^n \theta_i^2 + \frac{1}{6} \sum_{i=1}^n \theta_i^3 + \sum_{k=4}^{\infty} \frac{1}{k!} \left(\sum_{i=1}^n \theta_i^k \right)^{\frac{k}{4}} \\
 &= n - 1 + \sum_{i=1}^n \theta_i + \frac{1}{2} N_2 + \frac{1}{6} N_3 - N_4^{1/4} - \frac{1}{2} N_4^{1/2} - \frac{1}{6} N_4^{3/4} + \sum_{k=0}^{\infty} \frac{\sqrt[4]{N_4^k}}{k!} \\
 &= n - 1 + \gamma_c + \frac{2m + \gamma_c}{2} + \frac{N_3}{6} - N_4^{1/4} - \frac{N_4^{1/2}}{2} - \frac{N_4^{3/4}}{6} + e^{\sqrt[4]{N_4}}.
 \end{aligned}$$

Therefore, by Lemma 5 and Lemma 2, we get

$$EE_c(G) \leq n - 1 + \gamma_c + \frac{2m + \gamma_c}{2} + \frac{N_3}{6} - N_4^{1/4} - \frac{N_4^{1/2}}{2} - \frac{N_4^{3/4}}{6} + e^{\sqrt[4]{(\theta_n^2 + \theta_1^2)(2m + \gamma_c(G)) - n\theta_1^2\theta_n^2}}.$$

□

Theorem 4. Let G be a graph with n vertices and m edges. Then

$$EE_c(G) \geq \sqrt{n^2 + 4m + (n + 4) \gamma_c + \frac{4N_3}{3}}.$$

Proof. For any real x , one has

$$(17) \quad e^x \geq 1 + x + \frac{x^2}{2} + \frac{x^3}{3!}.$$

Since $e^x \geq 1 + x$, therefore, by this fact, Inequality (17) and the definition of the c -dominating Estrada index, we have

$$\begin{aligned}
 EE(G)^2 &= \left(\sum_{i=1}^n e^{\theta_i} \right)^2 \\
 &= \left(\sum_{i=1}^n e^{\theta_i} \right) (e^{\theta_1} + e^{\theta_n}) + \left(\sum_{i=1}^n e^{\theta_i} \right) (e^{\theta_2} + \dots + e^{\theta_{n-1}}) \\
 &\geq \sum_{i=1}^n e^{2\theta_i} + ne^{\theta_1 + \theta_n} + \left(\sum_{i=1}^n e^{\theta_i} \right) (e^{\theta_2} + \dots + e^{\theta_{n-1}}) \\
 &\geq \sum_{i=1}^n (1 + 2\theta_i + 2\theta_i^2 + \frac{8}{6}\theta_i^3) + ne^{\theta_1 + \theta_n} + \left(\sum_{i=1}^n e^{\theta_i} \right) (e^{\theta_2} + \dots + e^{\theta_{n-1}}) \\
 &= n + 2 \sum_{i=1}^n \theta_i + 2 \sum_{i=1}^n \theta_i^2 + \frac{8}{6} \sum_{i=1}^n \theta_i^3 + ne^{\theta_1 + \theta_n} + \left(\sum_{i=1}^n e^{\theta_i} \right) (e^{\theta_2} + \dots + e^{\theta_{n-1}}) \\
 &\geq n + 2\gamma_c + 2(2m + \gamma_c) + \frac{4N_3}{3} + n(1 + \theta_1 + \theta_n) + n((1 + \theta_2) + \dots + (1 + \theta_{n-1})) \\
 &\geq n^2 + 4m + (n + 4) \gamma_c + \frac{4N_3}{3},
 \end{aligned}$$

and this leads to the desired bound.

□

4. BOUND FOR THE c -DOMINATING ESTRADA INDEX INVOLVING THE c -DOMINATING ENERGY

In this section, we obtain lower and upper boundf for c -dominating Estrada index relates to c -dominating energy.

Theorem 5. *The c -dominating Estrada index $EE_c(G)$ and the c -dominating energy $E_c(G)$ satisfy the following inequality:*

$$(18) \quad EE_c(G) \geq \frac{1}{2}E_c(G)(e-1) + (n - n_+) + \gamma_c(G).$$

Proof. Since $e^x \geq 1 + x$, equality holds if and only if $x = 0$ and $e^x \geq ex$, equality holds if and only if $x = 1$, we have

$$\begin{aligned} EE_c(G) &= \sum_{i=1}^n e^{\theta_i} = \sum_{\theta_i > 0} e^{\theta_i} + \sum_{\theta_i \leq 0} e^{\theta_i} \\ &\geq \sum_{\theta_i > 0} e\theta_i + \sum_{\theta_i \leq 0} (1 + \theta_i) \\ &= e(\theta_1 + \theta_2 + \cdots + \theta_{n_+}) + (n - n_+) + (\theta_{n_++1} + \cdots + \theta_n) \\ &= (e-1)(\theta_1 + \theta_2 + \cdots + \theta_{n_+}) + (n - n_+) + \sum_{i=1}^n \theta_i \\ &= \frac{1}{2}E_c(G)(e-1) + (n - n_+) + \gamma_c(G). \end{aligned}$$

□

Theorem 6. *Let G be a graph with largest eigenvalue θ_1 and let p, τ and q be, respectively, the number of positive, zero and negative eigenvalues of G . Then*

$$(19) \quad EE_c(G) \geq \tau + pe^{\frac{E_c(G)+\gamma_c}{2p}} + qe^{\frac{\gamma_c-E_c(G)}{2q}}.$$

Proof. Let $\theta_1 \geq \cdots \geq \theta_p$ be the positive, and $\theta_{n-q+1}, \dots, \theta_n$ be the negative eigenvalues of G . By Equality (3), we have

$$(20) \quad \sum_{i=1}^p \theta_i + \sum_{i=n-q+1}^n \theta_i = \gamma_c.$$

And by the definition of the c -dominating energy, we have

$$(21) \quad E_c(G) = \sum_{i=1}^p \theta_i + \sum_{i=n-q+1}^n |\theta_i|.$$

By using Equalities (20) and (21), we have

$$\begin{aligned}
 E_c(G) &= \sum_{i=1}^p \theta_i + \sum_{i=n-q+1}^n |\theta_i| \\
 &= \gamma_c - \sum_{i=n-q+1}^n \theta_i + \sum_{i=n-q+1}^n |\theta_i| \\
 (22) \quad &= \gamma_c + 2 \sum_{i=n-q+1}^n |\theta_i| \\
 &= \gamma_c - 2 \sum_{i=n-q+1}^n \theta_i.
 \end{aligned}$$

Therefore, we get

$$(23) \quad \sum_{i=n-q+1}^n \theta_i = \frac{\gamma_c - E_c(G)}{2}.$$

With the same argument as above as well as using Enequality (22), we have

$$(24) \quad \sum_{i=p}^n \theta_i = \frac{E_c(G) + \gamma_c}{2}.$$

By the *arithmetic-geometric* mean inequality and Enequality (23), we have

$$(25) \quad \sum_{i=1}^p e^{\theta_i} \geq pe^{\frac{(\theta_1 + \dots + \theta_p)}{p}} = pe^{\frac{E_c(G) + \gamma_c}{2p}}.$$

Similarly and by using Enequality (24), we can write

$$(26) \quad \sum_{i=n-q+1}^n e^{\theta_i} \geq qe^{\frac{\gamma_c - E_c(G)}{2q}}.$$

For the zero eigenvalues, we have

$$\sum_{i=p+1}^{n-q} e^{\theta_i} = \tau.$$

So we obtain

$$EE_c(G) \geq \tau + pe^{\frac{E_c(G) + \gamma_c}{2p}} + qe^{\frac{\gamma_c - E_c(G)}{2q}}.$$

□

Theorem 7. Let G be a non-empty graph with n vertices and m edges. Then

$$EE_c(G) - E_c(G) \leq n - 1 - \sqrt{2m + \gamma_c(G) - 1} + e^{\sqrt{2m + \gamma_c(G) - 1}}.$$

Proof. By Inequality (16), we have

$$\begin{aligned}
EE_c(G) &\leq n + \sum_{k \geq 1} \frac{1}{k!} \sum_{i=1}^{n_+} (\theta_i)^k = n + \frac{E_c(G)}{2} + \sum_{i=1}^{n_+} \sum_{k \geq 2} \frac{(\theta_i)^k}{k!} \\
&< n + E_c(G) + \sum_{i=1}^{n_+} \sum_{k \geq 2} \frac{(\theta_i)^k}{k!} \\
&= n + E_c(G) + \sum_{k \geq 2} \frac{1}{k!} \left(\sum_{i=1}^{n_+} \theta_i \right)^k \\
&\leq n + E_c(G) + \sum_{k \geq 2} \frac{1}{k!} \left(\sum_{i=1}^{n_+} \theta_i^2 \right)^{\frac{k}{2}} \\
&\leq n + E_c(G) + \sum_{k \geq 2} \frac{1}{k!} (2m + \gamma_c - 1)^{\frac{k}{2}}.
\end{aligned}$$

Therefore, we have

$$EE_c(G) - E_c(G) \leq n - 1 - \sqrt{2m + \gamma_c(G) - 1} + e^{\sqrt{2m + \gamma_c(G) - 1}}.$$

□

Theorem 8. *Let G be a graph with n vertices. Then*

$$EE_c(G) \leq n - 1 + e^{E_c(G)},$$

and equality holds if and only if $G \cong \bar{K}_n$.

Proof. By definition of c -dominating Estrada index, we have

$$\begin{aligned}
EE_c(G) &\leq n + \sum_{i=1}^n \sum_{k \geq 1} \frac{|\theta_i|^k}{k!} \\
&= n + \sum_{k \geq 1} \frac{1}{k!} \left(\sum_{i=1}^n |\theta_i|^k \right) \\
&\leq n + \sum_{k \geq 1} \frac{1}{k!} \left(\sum_{i=1}^n |\theta_i| \right)^k \\
&= n + \sum_{k \geq 1} \frac{(E_c(G))^k}{k!}
\end{aligned}$$

which implies

$$EE_c(G) \leq n - 1 + e^{E_c(G)}.$$

It is not difficult to see that equality holds if and only if $G \cong \bar{K}_n$.

□

CONCLUDING REMARKS

In this paper, the c -dominating Estrada index of a graph is introduced and the c -dominating energy and the c -dominating Estrada index are studied. Moreover, we investigate the relations between the c -dominating Estrada index and the c -dominating energy.

ACKNOWLEDGMENT

We would like to thank the referee for their careful review of the paper and some helpful comments.

REFERENCES

- [1] N. Alawiah, N. J. Rad, A. Jahanbani, and H. Kamarulhaili, *New Upper Bounds on the Energy of a Graph*, MATCH Commun. Math. Comput. Chem. **79**, (2018), pp. 287–301.
- [2] G. Caporossi, D. Cvetković, I. Gutman, and P. Hansen, *Variable neighbourhood search for extremal graphs.2. Finding graphs with extremal energy*, J. Chem. Inf. Comput. Sci. **39** (1999), pp. 984–996.
- [3] K. CH. Das and I. Gutman, *Bounds for the energy of graphs*, Hacettepe Journal of Mathematics and Statistics, **45** (2016), pp. 695–703.
- [4] J. A. De la Peña and I. Gutman, J. Rada, *Estimating the Estrada Index*, Lin. Algebra Appl, **427** (2007), 70–76.
- [5] J. B. Diaz and F. T. Metcalf, *Stronger forms of a class of inequalities of G. Pólya - G. Szego, L. V. Kantorovich*, Bull. Amer. Math. Soc. **69** (1963), pp. 415–418.
- [6] E. Estrada and J. A. Rodríguez-Velázquez, *Subgraph Centrality in Complex Networks*, Phys. Rev. E, **71** (2005), 056103.
- [7] E. Estrada, *Characterization of 3D molecular structure*, Chem. Phys. Lett, **319** (2000), pp. 713–718.
- [8] E. Estrada, *Characterization of the folding degree of proteins*, Bioinformatics **18** (2002), pp. 697–704.
- [9] I. Gutman, *Bounds for total π -electron energy*, Chem. Phys. Lett. **24**(1974), pp. 283–285.
- [10] I. Gutman, *McClelland-type lower bound for total π -electron energy*, J. Chem. Soc. Faraday Trans. **86**(1990), pp. 3373–3375.
- [11] S. M. Hosamania, V. B. Awatib and R. M. Honmore, *On Graphs with equal Dominating and c -dominating energy*, arXiv preprint arXiv, **1811** 06269 (2018).
- [12] R. A. Horn and C. R. Johnson, *Matrix Analysis*, Cambridge Univ. Press, New York, 1985.
- [13] N. Jafari. Rad, A. Jahanbani, and D. A. Mojdeh, *Tetracyclic Graphs with Maximal Estrada Index*, Discrete Mathematics, Algorithms and Applications, **09** (2017), pp. 1750041:1–1750041:18.
- [14] N. Jafari. Rad, A. Jahanbani, and R. Hasni, *Pentacyclic Graphs with Maximal Estrada Index*, Ars Combin, **133** (2017), pp. 133–145.
- [15] N. Jafari. Rad, A. Jahanbani, and I. Gutman, *Zagreb Energy and Zagreb Estrada Index of Graphs*, MATCH Commun. Math. Comput. Chem, **79** (2018), pp. 371–386.
- [16] A. Jahanbani and J. R. Zambrano, *Koolen-Moulton-Type Upper Bounds on the Energy of a Graph*, MATCH Commun. Math. Comput. Chem, **83** (2020), pp. 497–518.
- [17] A. Jahanbani, *Upper bounds for the energy of graphs*, MATCH Commun. Math. Comput. Chem. **79** pp. 275–286.

- [18] A. Jahanbani, *Some new lower bounds for energy of graphs*, Applied Mathematics and Computation, **296** (2017), pp. 233–238.
- [19] A. Jahanbani, *Lower Bounds for the Energy of Graphs*, AKCE International Journal of Graphs and Combinatorics, **15** (2018), pp. 88–96.
- [20] A. Jahanbani, A. T. Karimi, and J. Rodriguez, *Results on the Estrada indices of Benzenoid Hydrocarbons Polycyclic Aromatic Compounds*. (2020), pp. 1–18.
- [21] A. Jahanbani, *On The Hermitian Estrada index of Mixed Graphs*, International Journal of Applied Graph Theory, **2** (2019), pp. 14–28.
- [22] A. Jahanbani, *Hermitian energy and Hermitian Estrada index of digraphs*, Asian-European Journal of Mathematics, **06** (2020), 2050116.
- [23] A. Jahanbani, *Albertson energy and Albertson Estrada index of graphs*, Journal of Linear and Topological Algebra (JLTA), **8** (2019), pp.11–24.
- [24] A. Jahanbani, *New Bounds for the harmonic energy and harmonic Estrada index of graphs*, Computer Science Journal of Moldova, **3** (2018), pp. 270–300.
- [25] X. Li, Y. Shi, and I. Gutman, *Graph Energy*, Springer, New York, 2012.
- [26] W. Lin, X. Guo, *Ordering trees by their largest eigenvalues*, Linear Algebra Appl. **418** (2006), pp.450–456.
- [27] M. A. Rashid, S. Ahmad, M. K. Siddiqui, A. Jahanbani, S. M. Sheikholeslami, and Zehui Shao, *New Bounds for the Estrada Index of Phenylenes Polycyclic Aromatic Compounds* (2020), pp. 1–17.
- [28] E. Sampathkumar and H. Walikar, *The connected domination number of a graph*, Journal of Mathematical and Physical Sciences, **13** (1979), pp. 607–613.

DEPARTMENT OF MATHEMATICS, AZARBAIJAN SHAHID MADANI UNIVERSITY, TABRIZ 51368, IRAN
Email address: Akbar.jahanbani92@gmail.com

DEPARTMENT OF MATHEMATICS, AZARBAIJAN SHAHID MADANI UNIVERSITY, TABRIZ 51368, IRAN
Email address: hajarshooshtary91@gmail.com